A Level Physics CIE

YOUR NOTES

2. Kinematics

CONTENTS

2.1 Equations of Motion
- 2.1.1 Displacement, Velocity & Acceleration
- 2.1.2 Motion Graphs
- 2.1.3 Area under a Velocity-Time Graph
- 2.1.4 Gradient of a Displacement-Time Graph
- 2.1.5 Gradient of a Velocity-Time Graph
- 2.1.6 Deriving Kinematic Equations
- 2.1.7 Solving Problems with Kinematic Equations
- 2.1.8 Acceleration of Free Fall Experiment
- 2.1.9 Projectile Motion

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

2.1 Equations of Motion

2.1.1 Displacement, Velocity & Acceleration

Defining Displacement, Velocity & Acceleration

Scalar quantities

- Remember scalar quantities only have a magnitude (size)
  - **Distance:** the total length between two points
  - **Speed:** the total distance travelled per unit of time

Vector quantities

- Remember vector quantities have both magnitude and direction
  - **Displacement:** the distance of an object from a fixed point in a specified direction
  - **Velocity:** the rate of change of displacement of an object
  - **Acceleration:** the rate of change of velocity of an object

Equations


SPEED AND VELOCITY ARE MEASURED IN METRES PER SECOND (ms⁻¹)
VELOCITY = CHANGE IN DISPLACEMENT / TIME
v = Δs/Δt
ACCELERATION = CHANGE IN VELOCITY / TIME
a = Δv/Δt
ACCELERATION IS MEASURED IN METRES PER SECOND EACH SECOND (ms⁻²)
IN PHYSICS, THE SYMBOL Δ MEANS "CHANGE"
Δs = CHANGE IN DISPLACEMENT
Δt = CHANGE IN TIME
Δv = CHANGE IN VELOCITY
Copyright © Save My Exams. All Rights Reserved

Equations linking displacement, velocity and acceleration

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Motion Graphs

- Three types of graph that can represent motion are **displacement-time** graphs, **velocity-time** graphs and **acceleration-time** graphs
- On a **displacement-time graph**...
  - **slope** equals **velocity**
  - the **y-intercept** equals the **initial displacement**
  - a **straight**(diagonal) line represents a **constant** velocity
  - a **curved** line represents an **acceleration**
  - a **positive slope** represents motion in the **positive direction**
  - a **negative slope** represents motion in the **negative direction**
  - a **zero** slope (horizontal line) represents a state of **rest**
  - the area under the curve is meaningless



Copyright © Save My Exams. All Rights Reserved

- On a **velocity-time graph**...
  - **slope** equals **acceleration**
  - the **y-intercept** equals the **initial velocity**
  - a **straight** line represents **uniform** acceleration
  - a **curved** line represents **non-uniform acceleration**
  - a **positive** slope represents an **increase** in **velocity** in the **positive direction**
  - a **negative** slope represents an **increase** in **velocity** in the **negative direction**
  - a **zero** slope (horizontal line) represents motion with **constant velocity**
  - the **area** under the curve equals the **displacement** or **distance** travelled

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES



VELOCITY–TIME GRAPH FOR CONSTANT VELOCITY



VELOCITY–TIME GRAPH FOR INCREASING VELOCITY



VELOCITY–TIME GRAPH FOR INCREASING ACCELERATION

Copyright © Save My Exams. All Rights Reserved

- On an **acceleration-time graph**...
  - slope is meaningless
  - the y-intercept equals the initial acceleration
  - a zero slope (horizontal line) represents an object undergoing constant acceleration
  - the area under the curve equals the change in velocity



ACCELERATION–TIME GRAPH FOR CONSTANT VELOCITY



ACCELERATION–TIME GRAPH FOR INCREASING VELOCITY



ACCELERATION–TIME GRAPH FOR INCREASING ACCELERATION

Copyright © Save My Exams. All Rights Reserved

How displacement, velocity and acceleration graphs relate to each other

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Area under a Velocity-Time Graph

- Velocity-time graphs show the speed and direction of an object in motion over a specific period of **time**
- The area under a velocity-time graph is equal to the **displacement** of a moving object

displacement = area under a velocity-time graph

Worked Example

The velocity-time graph of a vehicle travelling with uniform acceleration is shown in the diagram below.



Calculate the displacement of the vehicle at 40 s.

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

THE DISPLACEMENT IS EQUAL TO THE AREA UNDER A VELOCITY-TIME GRAPH



BASE = TIME = 40s

HEIGHT = VELOCITY = 105 km h^{-1}

CONVERT km h^{-1} TO km s^{-1}

$\frac{105}{60 \times 60} = 0.0292$ km s^{-1}

AREA OF A TRIANGLE = $\frac{1}{2}$ × BASE × HEIGHT

WORK OUT THE DISPLACEMENT

DISPLACEMENT = VELOCITY × TIME = $\frac{1}{2} \times 40 \times 0.0292 = 0.6$ km OR 600 m

Copyright © Save My Exams. All Rights Reserved

Exam Tip

Always check the values given on the y-axis of a motion graph - students often confuse displacement-time graphs and velocity-time graphs.The area under the graph can often be broken down into triangles, squares and rectangles, so make sure you are comfortable with calculating area!

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Gradient of a Displacement-Time Graph

- Displacement-time graphs show the changing position of an object in motion
- They also show whether an object is moving forwards (positive displacement) or backwards (negative displacement)
  - A negative gradient = a negative velocity (the object is moving backwards)
- The gradient (slope) of a displacement-time graph is equal to **velocity**
  - The greater the slope, the greater the velocity

Worked Example

A car driver sees a hazard ahead and applies the brakes to bring the car to rest.What does the displacement-time graph look like?



© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES


VELOCITY IS EQUAL TO THE GRADIENT OF THE DISPLACEMENT–TIME GRAPH
DISPLACEMENT IS INCREASING AT A CONSTANT RATE
THE GRADIENT OF THE DISPLACEMENT–TIME GRAPH IS DECREASING
S
0
t_1
t_2
t
V
VELOCITY IS CONSTANT
VELOCITY IS DECREASING AT A CONSTANT RATE
Copyright © Save My Exams. All Rights Reserved

Exam Tip

Don't forget that velocity is a vector quantity; it has a size and a direction. If velocity is initially positive and then becomes negative, then the object has changed direction.

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Gradient of a Velocity-Time Graph

- **Acceleration** is any change in the velocity of an object in a given time

$$acceleration = \frac{change\ in\ velocity}{time} = \frac{(v-u)}{t}$$

- As velocity is a vector quantity, this means that if the **speed** of an object **changes**, or its **direction changes**, then it is accelerating
  - An object that slows down tends to be described as 'decelerating'
- The gradient of a velocity-time graph is equal to **acceleration**

Worked Example

What does the velocity-time graph look like for this acceleration-time graph?



© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES


WHEN THE ACCELERATION REACHES A MAXIMUM, THE GRADIENT OF THE VELOCITY–TIME GRAPH STOPS INCREASING
ACCELERATION
0
TIME
ACCELERATION IS EQUAL TO THE GRADIENT OF A VELOCITY–TIME GRAPH
VELOCITY
0
TIME
WHEN THE ACCELERATION INCREASES THE GRADIENT OF THE VELOCITY–TIME GRAPH INCREASES
WHEN THE ACCELERATION DECREASES THE GRADIENT OF THE VELOCITY–TIME GRAPH DECREASES

Copyright © Save My Exams. All Rights Reserved

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

2.1.6 Deriving Kinematic Equations

Deriving Kinematic Equations of Motion

- The kinematic equations of motion are a set of four equations which can describe any object moving with **constant** acceleration
- They relate the five variables:
  - s = **displacement**
  - u = **initial velocity**
  - v = **final velocity**
  - a = **acceleration**
  - t = **time interval**
- It's important to know where these equations come from and how they are derived:





FROM THE GRADIENT WE CAN DEDUCE ACCELERATION IS EQUAL TO

- $\frac{\Delta y}{\Delta x} = \frac{\Delta v}{\Delta t} = \frac{(v-u)}{t}$
- $a = \frac{(v-u)}{t}$

MULTIPLY BOTH SIDES BY t

- $at = (v-u)$

REARRANGING LEADS TO

- $v = u + at$

Copyright © Save My Exams. All Rights Reserved

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

A graph showing how the velocity of an object varies with time

Deriving $s = \frac{(u + v)}{2} t$



THE OBJECT'S AVERAGE VELOCITY IS HALF-WAY BETWEEN u AND v:

$$\frac{(v + u)}{2}$$

DISPLACEMENT IS EQUAL TO AVERAGE VELOCITY × TIME SO:

$$s = \frac{(v + u)}{2} t$$

Copyright © Save My Exams. All Rights Reserved

The average velocity is halfway between u and v

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Deriving $s = ut + \frac{1}{2}at^2$



THE TWO TERMS ut AND 1/2at² MAKE UP THE AREA UNDER THE GRAPH

TAKING THE EQUATIONS WE DERIVED ABOVE

- $v = u + at$ (1)
- $s = \frac{(v + u)}{2}t$ (2)

SUBSTITUTING EQUATION (1) AND (2)

MULTIPLY EVERYTHING IN THE BRACKET BY t

- $s = \frac{(u + u + at)}{2}t$
- $s = \frac{2ut}{2} + \frac{at^2}{2}$

SEPARATE THE t AND t^2 TERMS

- $s = ut + \frac{1}{2}at^2$

Copyright © Save My Exams. All Rights Reserved

The two terms ut and ½at² make up the area under the graph

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Deriving $v^2 = u^2 + 2as$

TAKING THE EQUATIONS WE DERIVED ABOVE

- $v = u + at \longrightarrow t = \frac{v-u}{a}$ (1)

$s = \frac{(v+u)}{2}t$ (2)

SUBSTITUTING (1) INTO (2)

- $s = \frac{(v+u)}{2} \times \frac{(v-u)}{a}$
- $s = \frac{v^2-u^2}{2a}$ ← MULTIPLY BOTH SIDES BY 2a
- $v^2 = u^2 + 2as$

Copyright © Save My Exams. All Rights Reserved

This final equation can be derived from two of the others

SUMMARY

$v = u + at$

$s = ut + \frac{1}{2}at^2$

$s = \frac{(v+u)}{2}t$

$v^2 = u^2 + 2as$

Copyright © Save My Exams. All Rights Reserved

Summary of the four equations of uniformly accelerated motion

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Solving Problems with Kinematic Equations

- **Step 1:** Write out the variables that are given in the question, both known and unknown, and use the context of the question to deduce any quantities that aren't explicitly given
  - e.g. for vertical motion $a = \pm 9.81$ m s^{-2}, an object which starts or finishes at rest will have $u = 0$ or $v = 0$
- **Step 2:** Choose the equation which contains the quantities you have listed
  - e.g. the equation that links s, u, a and t is $s = ut + \frac{1}{2}at^2$
- **Step 3:** Convert any units to SI units and then insert the quantities into the equation and rearrange algebraically to determine the answer

Worked Example

The diagram shows an arrangement to stop trains that are travelling too fast. Trains



coming from the left travel at a speed of 50 ms^{-1}. At marker 1, the driver must apply the brakes so that the train decelerates uniformly in order to pass marker 2 at no more than 10 ms^{-1}.The train carries a detector that notes the times when the train passes each marker and will apply an emergency brake if the time between passing marker 1 and marker 2 is less than 20 s.How far from marker 2 should marker 1 be placed?

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

STEP 1

OUR KNOWN VARIABLES ARE

- $u = 50\ ms^{-1}$
- $v = 10\ ms^{-1}$
- $t = 20\ s$

AND WE ARE ASKED TO FIND DISTANCE, s.

STEP 2

SO THE EQUATION THAT LINKS u, v, t AND s IS

$$s = \frac{(u + v)}{2} t$$

STEP 3

NO REARRANGING IS REQUIRED SO WE SIMPLY PLUG IN THE VARIABLES:

$$s = \frac{(50 + 10)}{2} \times 20 = 30 \times 20 = 600\ m$$

Copyright © Save My Exams. All Rights Reserved

Exam Tip

- This is arguably the most important section of this topic, you can always be sure there will be one, or more, questions in the exam about solving problems with the kinematic equations
- The best way to master this section is to practice as many questions as possible

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Acceleration of Free Fall Experiment

- A common experiment to determine acceleration of a falling object which can be carried out in the lab

Apparatus

- Metre rule, ball bearing, electromagnet, electronic timer, trapdoor



Copyright © Save My Exams. All Rights Reserved

Apparatus used to measure g

Method

- When the current to the magnet switches off, the ball drops and the timer starts
- When the ball hits the trapdoor, the timer stops
- The reading on the timer indicates the time it takes for the ball to fall a distance, *h*
- This procedure is repeated several times for different values of *h*, in order to reduce random error
- The distance, *h*, can be measured using a metre rule as it would be preferable to use for distances between 20 cm – 1 m

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Analysing data

- To find g , use the same steps as in the problem solving section
- The known quantities are
  - Displacement $s = h$
  - Time taken = t
  - Initial velocity $u = 0$
  - Acceleration $a = g$
- The equation that links these quantities is
  - $s = ut + \frac{1}{2} at^2$
  - $h = \frac{1}{2} gt^2$
- Using this equation, deduce g from the gradient of the graph of h against t^2

Sources of error

- **Systematic error:** residue magnetism after the electromagnet is switched off may cause the time to be recorded as longer than it should be
- **Random error:** large uncertainty in distance from using a metre rule with a precision of 1mm, or from parallax error

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Projectile Motion

- The trajectory of an object undergoing projectile motion consists of a **vertical** component and a **horizontal** component
  - These need to be evaluated separately
- Some key terms to know, and how to calculate them, are:
  - **Time of flight:** how long the projectile is in the air
  - **Maximum height attained:** the height at which the projectile is momentarily at rest
  - **Range:** the horizontal distance travelled by the projectile



VERTICAL MOTION (↑)

INITIAL SPEED, $u = u\sin\theta$

ACCELERATION, $a = 9.81\ ms^{-2}$

DISPLACEMENT = 0

TIME OF FLIGHT

$u = u\sin\theta \quad v = 0 \quad a = -g \quad t = ?$

THE EQUATION THAT RELATES THESE QUANTITIES IS

$v = u + at$

$0 = u\sin\theta - gt$

$t = \frac{u\sin\theta}{g}$

$2t = \frac{2u\sin\theta}{g}$

IF THE TIME TO MAXIMUM HEIGHT IS t, THEN THE TIME OF FLIGHT IS 2t

MAXIMUM HEIGHT ATTAINED

$u = u\sin\theta \quad v = 0 \quad a = -g \quad H = ?$

THE EQUATION THAT RELATES THESE QUANTITIES IS

$v^2 = u^2 + 2as$

$0 = (u\sin\theta)^2 - 2gH$

$2gH = (u\sin\theta)^2$

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES


$H = \frac{(u\sin\theta)}{2g}$
HORIZONTAL MOTION (→)
INITIAL SPEED, $u = u\cos\theta$
ACCELERATION, $a = 0$
DISPLACEMENT = R
RANGE (R)
$u = u\cos\theta$ $t = \frac{2u\sin\theta}{g}$ $a = 0$ $R = ?$
THE EQUATION THAT RELATES THESE QUANTITIES IS
DISTANCE = SPEED × TIME
$R = (u\cos\theta)t$
$R = \frac{2u^2\sin\theta\cos\theta}{g}$
$R = \frac{u^2\sin 2\theta}{g}$
USING THE TRIG IDENTITY:
$2\sin\theta\cos\theta = \sin 2\theta$

Copyright © Save My Exams. All Rights Reserved

How to find the time of flight, maximum height and range

- **Remember:** the only force acting on the projectile, after it has been released, is **gravity**
- There are three possible scenarios for projectile motion:
  - **Vertical** projection
  - **Horizontal** projection
  - **Projection** at an **angle**
- Let's consider each in turn:

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Worked Example

To calculate vertical projection(free fall)

A science museum designed an experiment to show the fall of a feather in a vertical glass vacuum tube.

The time of fall from rest is 0.5 s.



Copyright © Save My Exams. All Rights Reserved

What it the length of the tube, L?

IN THIS PROBLEM, WE ONLY NEED TO CONSIDER VERTICAL MOTION. FIRST WE MUST LIST THE KNOWN VARIABLES.

$a = 9.81\ ms^{-2}$ $u = 0$ $t = 0.5\ s$ $L = ?$

THE EQUATION THAT LINKS THESE VARIABLES IS

$$s = ut + \frac{1}{2}at^2$$

$$L = \frac{1}{2}gt^2$$

$$L = \frac{1}{2} \times 9.81 \times 0.5^2 = 1.2\ m$$

Copyright © Save My Exams. All Rights Reserved

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Worked Example

To calculate horizontal projection

A motorcycle stunt-rider moving horizontally takes off from a point 1.25 m above the ground, landing 10 m away as shown.

What was the speed at take-off?



Copyright © Save My Exams. All Rights Reserved

IN THIS PROBLEM, WE NEED TO CONSIDER BOTH VERTICAL AND HORIZONTAL MOTION. LET'S CONSIDER THE VERTICAL MOTION FIRST. THE KNOWN VARIABLES ARE

$s = 1.25\ m$ $a = 9.81\ ms^{-2}$ $u = 0$ $t = ?$

THE EQUATION THAT LINKS THESE VARIABLES IS

$$s = ut + \frac{1}{2}at^2$$

$$s = \frac{1}{2}gt^2$$

$$t = \sqrt{\frac{2s}{g}}$$

$$t = \sqrt{\frac{2 \times 1.25}{9.81}} = 0.5s$$

NEXT LET'S CONSIDER THE HORIZONTAL MOTION. THE KNOWN VARIABLES ARE

$s = 10\ m$ $a = 0$ $t = 0.5\ s$ $u = ?$

SINCE THE ACCELERATION IS ZERO, WE CAN USE

$$\text{VELOCITY} = \frac{\text{DISPLACEMENT}}{\text{TIME}}$$

$$v = \frac{10}{0.5} = 20\ ms^{-1}$$

Copyright © Save My Exams. All Rights Reserved

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers

YOUR NOTES

Worked Example

To calculate projection at an angle

A ball is thrown from a point P with an initial velocity u of 12 m s^{-1} at 50° to the horizontal.

What is the value of the maximum height at Q?



Copyright © Save My Exams. All Rights Reserved

IN THIS PROBLEM, WE ONLY NEED TO CONSIDER VERTICAL MOTION UP TO THE POINT Q. FIRST WE MUST LIST THE KNOWN VARIABLES

$u = 12\sin(50)$ $a = -9.81\ ms^{-2}$ $v = 0$ $H = ?$

THE EQUATION THAT LINKS THESE VARIABLES IS

$$v^2 = u^2 + 2as$$

$$2as = v^2 - u^2$$

$$s = \frac{(v^2 - u^2)}{2a}$$

$$H = \frac{0 - (12\sin 50)^2}{2 \times (-9.81)}$$

$$H = \frac{(12\sin 50)^2}{19.62} = 4.3\ m$$

Copyright © Save My Exams. All Rights Reserved

Exam Tip

Make sure you don't make these common mistakes:

- Forgetting that deceleration is negative as the object rises
- Confusing the direction of sin θ and cos θ
- Not converting units (mm, cm, km etc.) to metres

© 2015-2021 Save My Exams, Ltd. · Revision Notes, Topic Questions, Past Papers